THE JOURNAL OF NUTRITION

OFFICIAL ORGAN OF THE AMERICAN INSTITUTE OF NUTRITION

GEORGE R. COWGILL, *Editor*
Yale Nutrition Laboratory, 333 Cedar Street
New Haven 11, Conn.

EDITORIAL BOARD

WILLIAM J. DARBY
D. M. HEGSTED
PEARL P. SWANSON
VERNON H. CHELDELIN
CARL V. MOORE
HAROLD H. WILLIAMS
ALEX BLACK
FLOYD S. DAFT
HARRY G. DAY
JAMES B. ALLISON
CARL A. BAUMANN
LEO C. NORRIS

VOLUME 47

MAY–AUGUST 1952

PUBLISHED MONTHLY BY
THE WISTAR INSTITUTE OF ANATOMY AND BIOLOGY
PHILADELPHIA, PENNSYLVANIA

CONTENTS

No. 1 MAY 1952

KNUT BREIREM. Biography of Oscar Kellner with frontispiece 3

MARY E. DUMM, ELAINE P. RALLI, HERBERT GERSHBERG AND BERTRAM LAKEN. The effects of diet, partial hepatectomy and growth-promoting factors on the composition of the rat liver. Three figures 11

SAMUEL M. GREENBERG WITH THE TECHNICAL ASSISTANCE OF LYDIA P. L. CHEN AND MARILYN A. BINGEMANN. The protective effect of dietary fat on immature rats fed thyroid 31

JOHN R. MCLEAN AND J. M. R. BEVERIDGE. Hepatic necrosis induced by dietary means. VI. The effect of varying the level and nature of protein and the level of fat on the development of liver necrosis 41

D. W. PETERSON, C. W. NICHOLS, JR. AND E. A. SHNEOUR WITH THE TECHNICAL ASSISTANCE OF H. GAFFEY, R. ROBBINS AND N. PEEK. Some relationships among dietary sterols, plasma and liver cholesterol levels, and atherosclerosis in chicks. One figure 57

MARY JONES HARP AND FLORENCE I. SCOULAR. Cobalt metabolism of young college women on self-selected diets 67

PENG-TUNG HSU AND G. F. COMBS. Effect of vitamin B_{12} and amino acid imbalances on growth and levels of certain blood constituents in the chick 73

HARRY P. BROQUIST, JOHN A. BROCKMAN, JR., MARVIN J. FAHRENBACH, E. L. R. STOKSTAD AND THOMAS H. JUKES. Comparative biological activity of leucovorin and pteroylglutamic acid. Three figures 93

K. SUVARNAKICH, G. V. MANN AND F. J. STARE. Riboflavin in human serum 105

IHSAN EL RAWI AND E. GEIGER. The growth-promoting effect of commercial strained meat and fish products investigated with infantile rats 119

H. H. MITCHELL AND JESSIE R. BEADLES. The determination of the protein requirement of the rat for maximum growth under conditions of restricted consumption of food. Two figures 133

BEULAH D. WESTERMAN, FAITH ROACH AND MIRIAM STONE. Improving the nutritive value of flour. V. The effect of the use of defatted wheat germ 147

No. 2 JUNE 1952

ALEX BLACK AND JOHN W. BRATZLER. The effects of a vitamin B_{12} supplement, vitamin B_{12} and streptomycin on the metabolism of the rat 159

G. W. ANDERSON, J. D. CUNNINGHAM AND S. J. SLINGER. Effect of protein level and penicillin on growth and intestinal flora of chickens. Five figures 175

JACK P. SAUNDERS, HAROLD R. SANDSTEAD, ROY E. BUTLER AND OLAF MICKELSEN. Relation of serum cholinesterase to nutritional status of adolescents 191

W. P. LEHRER, JR., A. C. WIESE AND P. R. MOORE. Biotin deficiency in suckling pigs. Three figures 203

GEORGE V. MANN, PATRICIA L. WATSON AND LUCILLE ADAMS. Primate nutrition. I. The Cebus monkey — Normal values. Four figures 213

GEORGE V. MANN, PATRICIA L. WATSON, ANNE MCNALLY AND JAMES GODDARD. Primate nutrition. II. Riboflavin deficiency in the Cebus monkey and its diagnosis. Six figures 225

KENNETH H. MADDY AND RAYMOND W. SWIFT. Growth and metabolism studies with rats fed rations containing purified amino acids. Two figures 243

E. A. KANE, W. C. JACOBSON AND L. A. MOORE. Diurnal variation in the excretion of chromium oxide and lignin. One figure 263

HERBERT P. SARETT. Effect of oral administration of streptomycin on urinary excretion of B vitamins in man 275

BENJAMIN H. ERSHOFF. Increased survival of liver-fed rats administered multiple sublethal doses of x-irradiation 289

GERALD LITWACK, J. N. WILLIAMS, JR., LILLIAN CHEN AND C. A. ELVEHJEM. A study of the relationship of liver xanthine oxidase to quality of dietary protein. Three figures 299

No. 3 JULY 1952

ALICE M. VIOLANTE, ROBERT J. SIRNY AND C. A. ELVEHJEM. The aspartic acid, glutamic acid, proline and tyrosine content of meat 307

MARGUERITE A. CONSTANT AND PAUL H. PHILLIPS. The occurrence of a calcinosis syndrome in the cotton rat. I. The effect of diet on the ash content of the heart. One figure 317

MARGUERITE A. CONSTANT, PAUL H. PHILLIPS AND D. MURRAY ANGEVINE. The occurrence of a calcinosis syndrome in the cotton rat. II. Pathology. Eleven figures 327

BESSIE L. DAVEY, MEI-LING WU AND CLARA A. STORVICK. With the technical assistance of Katherine Ding and Betty Jo Graves. Daily determination of plasma, serum and white cell-platelet ascorbic acid in relationship to the excretion of ascorbic and homogentisic acids by adults maintained on a controlled diet. One figure 341

WILLIAM T. BEHER, EVELYN M. CRIGGER AND O. H. GAEBLER. The effect of testosterone propionate on niacin metabolism in rats 353

H. D. ALEXANDER AND R. W. ENGEL. The importance of choline in the prevention of nutritional edema in rats fed low-protein diets. Three figures 361

JAMES W. BOTHWELL AND J. N. WILLIAMS, JR. A study of lysine deficiency and nitrogen balance under ad libitum and force-feeding conditions. One figure 375

E. KYLE BACON, SVEN LASSEN, SAMUEL M. GREENBERG, JOHN W. MEHL AND HARRY J. DEUEL, JR. The influence of ingested mineral oil upon the development of an essential fatty acid deficiency in the rat. Five figures 383

RALPH HOAGLAND, G. G. SNIDER AND C. E. SWIFT. Nutritive value of lard as affected by the proportion of fat in the diet 399

R. M. FORBES AND W. T. HAINES. The riboflavin requirement of the baby pig. I. At an environmental temperature of 85°F. and 70% relative humidity. Three figures 411

FRANCES A. JOHNSTON AND THELMA J. MCMILLAN. With the technical assistance of Gladys Derby Falconer and Erica Evans. The amount of nitrogen retained by 6 young women on an intake of approximately 70 gm of protein a day 425

E. S. NASSET AND ANN M. SILICIANO. Nitrogen balance of adult rats fed diets low in L-, DL- and D-phenylalanine. One figure 437

L. M. DANSKY AND F. W. HILL. Application of the chromic oxide indicator method to balance studies with growing chickens. Two figures 449

D. V. CATRON, D. RICHARDSON, L. A. UNDERKOFLER, H. M. MADDOCK AND W. C. FRIEDLAND. Vitamin B_{12} requirement of weanling pigs. II. Performance on low levels of vitamin B_{12} and requirement for optimum growth 461

No. 4 AUGUST 1952

R. A. SALANDER AND A. R. PATTON. Free amino acids in chick livers as altered by vitamin B_{12} and fasting 469

W. H. SEBRELL, JR. AND ERNEST G. MCDANIEL. Amino acids in the production of blood constituents in rats 477

WILLIAM H. PFANDER. The effects of iodinated casein, thiouracil and cortisone on the ascorbic acid requirement of the male guinea pig. Seven figures 487

W. H. PFANDER AND H. H. MITCHELL. The ascorbic acid requirement of the guinea pig when adrenal weight and odontoblast height are used as criteria. Five figures 503

CHARLES H. BARROWS, ERNESTINE BECKER MCCOLLUM AND BACON F. CHOW. The urinary excretion of vitamin B_{12} following subcutaneous administration to young and adult rats with different reserves. One figure 525

J. P. LAMBOOY AND H. V. APOSHIAN. The biological activity of diethyl riboflavin. Seven figures 539

G. P. LOFGREEN, MAX KLEIBER AND A. H. SMITH. The excretion of injected P^{32} into the gastrointestinal tract of the young calf. Three figures ... 561

G. P. LOFGREEN, MAX KLEIBER AND J. R. LUICK. The metabolic fecal phosphorus excretion of the young calf. Two figures 571

HAROLD H. DRAPER, MARIAN F. JAMES AND B. CONNOR JOHNSON. Tri-o-cresyl phosphate as a vitamin E antagonist for the rat and lamb. Four figures 583

HAROLD E. SCHEID, MONA M. ANDREWS AND B. S. SCHWEIGERT. Comparison of methods for determination of the vitamin B_{12} potency of meats. One figure 601

I. ZIPKIN AND F. J. MCCLURE. Deposition of fluorine in the bones and teeth of the growing rat. Four figures 611

J. E. CAMPBELL AND P. H. PHILLIPS. Reproduction studies with dogs. I. Effect of a corn-soybean meal-alfalfa diet. One figure 621

F. H. KRATZER, D. E. WILLIAMS AND BLANCHE MARSHALL. The requirement for isoleucine and the activities of its isomers for the growth of turkey poults. One figure 631

INDEX

AD libitum and force-feeding conditions, study of lysine deficiency and nitrogen balance under, 375.
ADAMS, LUCILLE. See Mann, G. V., 213.
Adolescents (see Man).
Adrenal weight and odontoblast heights as criteria for ascorbic acid requirement of guinea pig, 503.
Adult rats (see Rats, adult).
Adults (see Man).
ALEXANDER, H. D., AND R. W. ENGEL. The importance of choline in the prevention of nutritional edema in rats fed low-protein diets, 361.
Alfalfa-corn-soybean meal diet, effect of. Reproduction studies with dogs. I., 621.
Amino acid(s), free, in chick livers as altered by vitamin B_{12} and fasting, 469.
—— imbalances and vitamin B_{12}, effect on growth and levels of certain blood constituents in chick, 73.
—— in production of blood constituents in rats, 477.
—— purified, growth and metabolism studies with rats fed rations containing, 243.
ANDERSON, G. W., J. D. CUNNINGHAM AND S. J. SLINGER. Effect of protein level and penicillin on growth and intestinal flora of chickens, 175.
ANDREWS, MONA M. See Scheid, H. E., 601.
ANGEVINE, D. MURRAY. See Constant, M. A., 327.
Antagonist (vitamin E) for rat and lamb, tri-o-cresyl phosphate as a, 583.
APOSHIAN, H. V. See Lambooy, J. P., 539.
Application of chromic oxide indicator method to balance studies with growing chickens, 449.
Ascorbic acid (plasma, serum and white cell-platelet), daily determination of, in relation to excretion of ascorbic and homogentisic acids by adults maintained on controlled diet, 341.
—— requirement of guinea pig when adrenal weight and odontoblast height are used as criteria, 503.
—— —— —— male guinea pig, effects of iodinated casein, thiouracil and cortisone on, 487.
— and homogentisic acids, excretion by adults maintained on controlled diet, daily determination of plasma, serum and white cell-platelet ascorbic acid in relation to, 341.
Ash content of heart, effect of diet on. Occurrence of calcinosis syndrome in cotton rat. I., 317.
Aspartic acid content of meat, 307.
Atherosclerosis, dietary sterols, and plasma and liver cholesterol levels, some relationships among, in chicks, 57.

B VITAMINS (see Vitamins, B).
Baby pig (see Pig, baby).
BACON, E. K., S. LASSEN, S. M. GREENBERG, J. W. MEHL AND H. J. DEUEL, JR. The influence of ingested mineral oil upon the development of an essential fatty acid deficiency in the rat, 383.
Balance (nitrogen) and lysine deficiency, study of, under ad libitum and force-feeding conditions, 375.
—— of adult rats fed diets low in L-, DL- and D-phenylalanine, 437.
— studies with growing chickens, application of chromic oxide indicator method to, 449.
BARROWS, C. H., E. B. MCCOLLUM AND B. F. CHOW. The urinary excretion of vitamin B_{12} following subcutaneous administration to young and adult rats with different reserves, 525.
BEADLES, JESSIE R. See Mitchell, H. H., 133.
BEHLEE, W. T., E. M. GRIGGER AND O. H. GAEBLER. The effect of testosterone propionate on niacin metabolism in rats, 353.
BEVERIDGE, JAMES M. R. See McLean, J. R., 41.
BINGEMANN, MARILYN A. See Greenberg, S. M., 31.
Biological activity, comparative, of leucovorin and pteroylglutamic acid, 93.
—— of diethyl riboflavin, 539.
Biotin deficiency in suckling pigs, 203.
BLACK, A., AND J. W. BRATZLER. The effects of a vitamin B_{12} supplement, vitamin B_{12} and streptomycin on the metabolism of the rat, 159.
Blood constituents, certain, in chick, effect of vitamin B_{12} and amino acid imbalances on levels of, 73.
—— in rats, amino acids in production of, 477.
Bones and teeth of growing rat, deposition of fluorine in, 611.
BOTHWELL, J. W., AND J. N. WILLIAMS, JR. A study of lysine deficiency and nitrogen balance under ad libitum and force-feeding conditions, 375.
BRATZLER, JOHN W. See Black, A., 159.
BREIREM, K. Oscar Kellner (May 13, 1851–September 22, 1911), 3.
BROCKMAN, JOHN A., JR. See Broquist, H. P., 93.
BROQUIST, H. P., J. A. BROCKMAN, JR., M. J. FAHRENBACH, E. L. R. STOCKSTAD AND T. H. JUKES. Comparative biological activity of leucovorin and pteroylglutamic acid, 93.
BUTLER, ROY E. See Saunders, J. P., 191.

CALCINOSIS syndrome, occurrence in cotton rat. I. Effect of diet on ash content of heart, 317; II. Pathology, 327.
Calf, young, excretion of injected P^{32} into gastrointestinal tract of, 561.
—— metabolic fecal phosphorus excretion of, 571.
CAMPBELL, J. E., AND P. H. PHILLIPS. Reproduction studies with dogs. I. Effect of a corn-soybean meal-alfalfa diet, 621.
Casein, iodinated, effect on ascorbic acid requirement of male guinea pig, 487.
CATRON, D. V., D. RICHARDSON, L. A. UNDERKOFLER, H. M. MADDOCK AND W. C. FRIEDLAND. Vitamin B_{12} requirement of weanling pigs. II. Performance on low levels of vitamin B_{12} and requirement for optimum growth, 461.

Cebus monkey, normal values. Primate nutrition. I., 213.
—— riboflavin deficiency in, and its diagnosis, 225.
CHEN, LILLIAN. See Litwack, G., 299.
— LYDIA PI-LING. See Greenberg, S. M., 31.
Chick(s), effect of vitamin B_{12} and amino acid imbalances on growth and levels of certain block constituents in, 73.
— livers, free amino acids in, as altered by vitamin B_{12} and fasting, 469.
— some relationships among dietary sterols, plasma and liver cholesterol levels, and atherosclerosis in, 57.
Chickens, effect of protein level and penicillin on growth and intestinal flora of, 175.
— growing, application of chromic oxide indicator method to balance studies with, 449.
Cholesterol levels (plasma and liver), dietary sterols, and atherosclerosis, some relationships among, in chicks, 57.
Choline, importance in prevention of nutritional edema in rats fed low-protein diets, 361.
Cholinesterase, serum, relation to nutritional status of adolescents, 191.
CHOW, BACON F. See Barrows, C. H., 525.
Chromic oxide indicator method, application to balance studies with growing chickens, 449.
Chromium oxide and lignin, diurnal variation in excretion of, 263.
Cobalt metabolism (see Metabolism, cobalt).
College women (see Women, college).
COMBS, GERALD F. See Hsu, P.-T., 73.
Commercial strained meat and fish products, growth-promoting effect of, investigated with infantile rats, 119.
CONSTANT, M. A., AND P. H. PHILLIPS. The occurrence of a calcinosis syndrome in the cotton rat. I. The effect of diet on the ash content of the heart, 317.
—— —— P. H. PHILLIPS AND D. M. ANGEVINE. The occurrence of a calcinosis syndrome in the cotton rat. II. Pathology, 327.
Corn-soybean meal-alfalfa diet, effect of. Reproduction studies with dogs. I., 621.
Cortisone, effect on ascorbic acid requirement of male guinea pig, 487.
Cotton rat, occurrence of calcinosis syndrome in. I., 317; II., 327.
CRIGGER, EVELYN M. See Beher, W. T., 353.
Criteria of ascorbic acid requirement of guinea pig, adrenal weight and odontoblast height used as, 503.
CUNNINGHAM, JOHN D. See Anderson, G. W., 175.

DANSKY, L. M., AND F. W. HILL. Application of the chromic oxide indicator method to balance studies with growing chickens, 449.
DAVEY, B. L., M.-L. WU AND C. A. STORVICK. With the technical assistance of K. Ding and B. J. Graves. Daily determination of plasma, serum and white cell-platelet ascorbic acid in relation to the excretion of ascorbic and homogentisic acids by adults maintained on a controlled diet, 341.
Defatted wheat germ, effect of use of. Improving the nutritive value of flour. V., 147.
DEUEL, HARRY J., JR. See Bacon, E. K., 383.
Diagnosis of riboflavin deficiency in Cebus monkey, 225.
Diet(s), controlled, daily determination of plasma, serum and white cell-platelet ascorbic acid in relation to excretion of ascorbic and homogentisic acids by adults maintained on, 341.
— corn-soybean meal-alfalfa, effect of. Reproduction studies with dogs. I., 621.
— effect on ash content of heart. Occurrence of calcinosis syndrome in cotton rat. I., 317.
—— —— composition of rat liver, 11.
— low in L-, DL- and D-phenylalanine, nitrogen balance of adult rats fed, 437.
— low-protein, importance of choline in prevention of nutritional edema in rats fed, 361.
— nutritive value of lard as affected by proportion of fat in, 399.
— self-selected, cobalt metabolism of young college women on, 67.
Dietary fat, protective effect on immature rats fed thyroid, 31.
— means, hepatic necrosis induced by. VI., 41.
— protein, relationship of liver xanthine oxidase to quality of, 299.
— sterols, plasma and liver cholesterol levels, and atherosclerosis, some relationships among, in chicks, 57.
Diethyl riboflavin, biological activity of, 539.
DING, KATHERINE. See Davey, B. L., 341.
Diurnal variation in excretion of chromium oxide and lignin, 263.
Dogs, reproduction studies with. I., 621.
Doses, multiple sublethal, of x-irradiation, increased survival of liver-fed rats administered, 289.
DRAPER, H. H., M. F. JAMES AND B. C. JOHNSON. Tri-o-cresyl phosphate as a vitamin E antagonist for the rat and lamb, 583.
DUMM, M. E., E. P. RALLI, H. GERSHBERG AND B. LAKEN. The effects of diet, partial hepatectomy and growth-promoting factors on the composition of the rat liver, 11.

EDEMA, nutritional, in rats fed low-protein diets, importance of choline in prevention of, 361.
EL RAWI, I., AND E. GEIGER. The growth-promoting effect of commercial strained meat and fish products investigated with infantile rats, 119.
ELVEHJEM, CONRAD A. See Litwack, G., 299; Violante, A. M., 307.
ENGEL, R. W. See Alexander, H. D., 361.
ERSHOFF, B. H. Increased survival of liver-fed rats administered multiple sublethal doses of x-irradiation, 289.
EVANS, ERICA. See Johnston, F. A., 425.
Excretion, metabolic fecal phosphorus, of young calf, 571.
— of ascorbic and homogentisic acids by adults maintained on controlled diet, daily determination of plasma, serum and white cell-platelet ascorbic acid in relation to, 341.
—— chromium oxide and lignin, diurnal variation in, 263.
—— injected P^{32} into gastrointestinal tract of young calf, 561.
— urinary (see Urinary excretion).

FAHRENBACH, MARVIN J. See Broquist, H. P., 93.
FALCONER, GLADYS D. See Johnston, F. A., 425.
Fasting, free amino acids in chick livers as altered by, 469.
Fat, dietary, protective effect on immature rats fed thyroid, 31.
— effect of varying level of, and level and nature of protein, on development of liver necrosis, 41.
— in diet, nutritive value of lard as affected by proportion of, 399.
Fatty acid (essential) deficiency in rat, influence of ingested mineral oil upon development of, 383.
Fecal phosphorus (metabolic) excretion of young calf, 571.
Fish and meat products (commercial strained), growth-promoting effect of, investigated with infantile rats, 119.
Flora, intestinal (see Intestinal flora).
Flour, improving the nutritive value of. V., 147.

Fluorine, deposition in bones and teeth of growing rat, 611.
Food, determination of protein requirement of rat for maximum growth under conditions of restricted consumption of, 133.
FORBES, R. M., AND W. T. HAINES. The riboflavin requirement of the baby pig. I. At an environmental temperature of 85°F. and 70% relative humidity, 411.
Force-feeding and ad libitum conditions, study of lysine deficiency and nitrogen balance under, 375.
Free amino acids in chick livers as altered by vitamin B_{12} and fasting, 469.
FRIEDLAND, W. C. See Catron, D. V., 461.

GAEBLER, O. H. See Beher, W. T., 353.
GAFFEY, HAZEL. See Peterson, D. W., 57.
Gastrointestinal tract of young calf, excretion of injected P^{32} into, 561.
GEIGER, ERNEST. See El Rawi, I., 119.
Germ, defatted wheat, effect of use of. Improving the nutritive value of flour. V., 147.
GERSHBERG, HERBERT. See Dumm, M. E., 11.
Glutamic acid content of meat, 307.
GODDARD, JAMES. See Mann, G. V., 225.
GRAVES, BETTY JO. See Davey, B. L., 341.
GREENBERG, S. M. With the technical assistance of L. P.-L. Chen, and M. A. Bingemann. The protective effect of dietary fat on immature rats fed thyroid, 31.
— — — See Bacon, E. K., 383.
Growing chickens (see Chickens, growing).
— rat (see Rat, growing).
Growth and metabolism studies with rats fed rations containing purified amino acids, 243.
— maximum, under conditions of restricted consumption of food, determination of protein requirement of rat for, 133.
— of chick, effect of vitamin B_{12} and amino acid imbalances on, 73.
— of chicks, effect of protein level and penicillin on, 175.
— of turkey poults, requirement for isoleucine and activities of its isomers for, 631.
— optimum, vitamin B_{12} requirement of weanling pigs for, 461.
Growth-promoting effect of commercial strained meat and fish products investigated with infantile rats, 119.
— — factors, effect on composition of rat liver, 11.
Guinea pig, ascorbic acid requirement of, when adrenal weight and odontoblast height are used as criteria, 503.
— — male, effects of iodinated casein, thiouracil and cortisone on ascorbic acid requirement of, 487.

HAINES, W. T. See Forbes, R. M., 411.
HARP, M. J., AND F. I. SCOULAR. Cobalt metabolism of young college women on self-selected diets, 67.
Heart, effect of diet on ash content of. Occurrence of calcinosis syndrome in cotton rat, I., 317.
Height (odontoblast) and adrenal weight as criteria of ascorbic acid requirement of guinea pig, 503.
Hepatectomy, effect of partial, on composition of rat liver, 11.
Hepatic necrosis induced by dietary means. VI. The effect of varying the level and nature of protein and the level of fat on the development of liver necrosis, 41.
HILL, F. W. See Dansky, L. M., 449.
HOAGLAND, R., G. G. SNIDER AND C. E. SWIFT. Nutritive value of lard as affected by the proportion of fat in the diet, 399.
Homogentisic and ascorbic acids, excretion by adults maintained on controlled diet, daily determination of plasma, serum and white cell-platelet ascorbic acid in relation to, 341.
HSU, P.-T., AND G. F. COMBS. Effect of vitamin B_{12} and amino acid imbalances on growth and levels of certain blood constituents in the chick, 73.
Human serum, riboflavin in, 105.
Humidity, relative, of 70% and environmental temperature of 85°F., riboflavin requirement of baby pig at, 411.

IMBALANCES (amino acid) and vitamin B_{12}, effect on growth and levels of certain blood constituents in chick, 73.
Immature rats (see Rats, immature).
Indicator method (chromic oxide), application to balance studies with growing chickens, 449.
Infantile rats (see Rats, infantile).
Intestinal flora of chickens, effect of protein level and penicillin on, 175.
Iodinated casein, effect on ascorbic acid requirement of male guinea pig, 487.
Isoleucine, requirement for, and activities of its isomers for growth of turkey poults, 631.

JACOBSON, WESLEY C. See Kane, E. A., 263.
JAMES, MARIAN F. See Draper, H. H., 583.
JOHNSON, B. CONNOR. See Draper, H. H., 583.
JOHNSTON, F. A., AND T. J. MCMILLAN. With the technical assistance of G. D. Falconer and E. Evans. The amount of nitrogen retained by 6 young women on an intake of approximately 70 gm of protein a day, 425.
JUKES, THOMAS H. See Broquist, H. P., 93.

KANE, E. A., W. C. JACOBSON AND L. A. MOORE. Diurnal variation in the excretion of chromium oxide and lignin, 263.
KELLNER, OSCAR (May 13, 1851–September 22, 1911), 3.
KLEIBER, MAX. See Lofgreen, G. P., 561, 571.
KRATZER, F. H., D. E. WILLIAMS AND B. MARSHALL. The requirement for isoleucine and the activities of its isomers for the growth of turkey poults, 631.

LAKEN, BERTRAM. See Dumm, M. E., 11.
Lamb, tri-o-cresyl phosphate as a vitamin E antagonist for, 583.
LAMBOOY, J. P., AND H. V. APOSHIAN. The biological activity of diethyl riboflavin, 539.
Lard, nutritive value as affected by proportion of fat in diet, 399.
LASSEN, SVEN. See Bacon, E. K., 383.
LEHRER, W. P., JR., A. C. WIESE AND P. R. MOORE. Biotin deficiency in suckling pigs, 203.
Leucovorin and pteroylglutamic acid, comparative biological activity of, 93.
Lignin and chromium oxide, diurnal variation in excretion of, 263.
LITWACK, G., J. N. WILLIAMS, JR., L. CHEN AND C. A. ELVEHJEM. A study of the relationship of liver xanthine oxidase to quality of dietary protein, 299.
Liver(s) and plasma cholesterol levels, dietary sterols, and atherosclerosis, some relationship among, in chicks, 57.
— chick, free amino acids in, as altered by vitamin B_{12} and fasting, 469.
— necrosis, effect of varying level and nature of protein and level of fat on development of, 41.

— rat, effects of diet, partial hepatectomy and growth-promoting factors on composition of, 11.
— xanthine oxidase, relationship to quality of dietary protein, 299.
Liver-fed rats administered multiple sublethal doses of x-irradiation, increased survival of, 289.
LOFGREEN, G. P., M. KLEIBER AND A. H. SMITH. The excretion of injected P^{32} into the gastrointestinal tract of the young calf, 561.
— — — — — — — — J. R. LUICK. The metabolic fecal phosphorus excretion of the young calf, 571.
LUICK, J. R. See Lofgreen, G. P., 571.
Lysine deficiency and nitrogen balance, study of, under ad libitum and force-feeding conditions, 375.

MADDOCK, H. M. See Catron, D. V., 461.
MADDY, K. H., AND R. W. SWIFT. Growth and metabolism studies with rats fed rations containing purified amino acids, 243.
Male guinea pig (see Guinea pig, male).
Man, effect of oral administration of streptomycin on urinary excretion of B vitamins in, 275.
— maintained on controlled diet, daily determination of plasma, serum and white cell-platelet ascorbic acid in relation to excretion of ascorbic and homogentisic acids by, 341.
— relation of serum cholinesterase to nutritional status of, 191.
MANN, G. V., P. L. WATSON AND L. ADAMS. Primate nutrition. I. The Cebus monkey — normal values, 213.
— — — — — — — — ANNE MCNALLY AND J. GODDARD. Primate nutrition. II. Riboflavin deficiency in the Cebus monkey and its diagnosis, 225.
— — — — See Suvarnakich, K., 105.
MARSHALL, BLANCHE. See Kratzer, F. H., 631.
MCCLURE, F. J. See Zipkin, I., 611.
MCCOLLUM, ERNESTINE B. See Barrows, C. H., 525.
MCDANIEL, ERNEST G. See Sebrell, W. H., Jr., 477.
MCLEAN, J. R., AND J. M. R. BEVERIDGE. Hepatic necrosis induced by dietary means. VI. The effect of varying the level and nature of protein and the level of fat on the development of liver necrosis, 41.
MCMILLAN, THELMA J. See Johnston, F. A., 425.
MCNALLY, ANNE. See Mann, G. V., 225.
Meat(s) and fish products (commercial strained), growth-promoting effect of, investigated with infantile rats, 119.
— aspartic acid, glutamic acid, proline and tyrosine content of, 307.
— comparison of methods for determination of vitamin B_{12} potency of, 601.
MEHL, JOHN W. See Bacon, E. K., 383.
Metabolic fecal phosphorus excretion of young calf, 571.
Metabolism and growth studies with rats fed rations containing purified amino acids, 243.
— cobalt, of young college women on self-selected diets, 67.
— niacin, in rats, effect of testosterone propionate on, 353.
— of rats, effects of vitamin B_{12} supplement, vitamin B_{12} and streptomycin on, 159.
Methods for determination of vitamin B_{12} potency of meats, comparison of, 601.
MICKELSEN, OLAF. See Saunders, J. P., 191.
Mineral oil, influence of ingested, upon development of an essential fatty acid deficiency in rat, 383.
MITCHELL, H. H., AND J. R. BEADLES. The determination of the protein requirement of the rat for maximum growth under conditions of restricted consumption of food, 133.
— — — See Pfander, W. H., 503.
Monkey, Cebus, normal values. Primate nutrition. I., 213.
— — riboflavin deficiency in, and its diagnosis, 225.
MOORE, LANE A. See Kane, E. A., 263.
— PETER R. See Lehrer, W. P., Jr., 203.
Multiple sublethal doses of x-irradiation, increased survival of liver-fed rats administered, 289.

NASSET, E. S., AND A. M. SILICIANO. Nitrogen balance of adult rats fed diets low in L-, DL- and D-phenylalanine, 437.
Necrosis, hepatic, induced by dietary means. VI., 41.
Niacin metabolism (see Metabolism, niacin).
NICHOLS, CLARENCE W., JR. See Peterson, D. W., 57.
Nitrogen, amount retained by 6 young women on intake of approximately 70 gm protein a day, 425.
— balance and lysine deficiency, study of, under ad libitum and force-feeding conditions, 375.
— — of adult rats fed diets low in L-, DL- and D-phenylalanine, 437.
Nutrition, primate. I., 213; II., 225.
Nutritional edema in rats fed low-protein diets, importance of choline in prevention of, 361.
— status of adolescents, relation of serum cholinesterase to, 191.
Nutritive value of flour, improving the. V. The effect of the use of defatted wheat germ, 147.
— — — lard as affected by proportion of fat in diet, 399.

ODONTOBLAST height and adrenal weight as criteria of ascorbic acid requirement of guinea pig, 503.
Oral administration of streptomycin, effect on urinary excretion of B vitamins in man, 275.
Oxidase (liver xanthine), relationship to quality of dietary protein, 299.
Oxide, chromic (see Chromic oxide).
— (chromium) and lignin, diurnal variation in excretion of, 263.

P^{32}, excretion of injected, into gastrointestinal tract of young calf, 561.
Pathology of calcinosis syndrome in cotton rat, 327.
PATTON, A. R. See Salander, R. C., 469.
PEEK, NEAL. See Peterson, D. W., 57.
Penicillin and protein level, effect on growth and intestinal flora of chickens, 175.
PETERSON, D. W., C. W. NICHOLS, JR. AND E. A. SHNEOUR. With the technical assistance of H. Gaffey, R. Robbins and N. Peek. Some relationships among dietary sterols, plasma and liver cholesterol levels, and atherosclerosis in chicks, 57.
PFANDER, W. H. The effects of iodinated casein, thiouracil and cortisone on the ascorbic acid requirement of the male guinea pig, 487.
— — — AND H. H. MITCHELL. The ascorbic acid requirement of the guinea pig when adrenal weight and odontoblast height are used as criteria, 503.
Phenylalanine (L-, DL- AND D-), nitrogen balance of adult rats fed diets low in, 437.
PHILLIPS, P. H. See Campbell, J. E., 621; Constant, M. A., 317, 327.
Phosphate (tri-o-cresyl) as a vitamin E antagonist for rat and lamb, 583.

Phosphorus (metabolic fecal) excretion of young calf, 571.
Pig(s), baby, riboflavin requirement of. I., 411.
— guinea (see Guinea pig).
— suckling, biotin deficiency in, 203.
— weanling, vitamin B_{12} requirement of, II., 461.
Plasma and liver cholesterol levels, dietary sterols, and atherosclerosis, some relationships among, in chicks, 57.
— serum and white cell-platelet ascorbic acid, daily determination of, in relation to excretion of ascorbic and homogentisic acids by adults maintained on controlled diet, 341.
Platelet-white cell, plasma and serum ascorbic acid, daily determination of, in relation to excretion of ascorbic and homogentisic acids by adults maintained on controlled diet, 341.
Poults, turkey (see Turkey poults).
Primate nutrition. I. Cebus monkey — normal values, 213; II. Riboflavin deficiency in Cebus monkey and its diagnosis, 225.
Proline content of meat, 307.
Propionate, testosterone (see Testosterone propionate).
Protective effect of dietary fat on immature rats fed thyroid, 31.
Protein, amount of nitrogen retained by 6 young women on intake of approximately 70 gm a day of, 425.
— dietary, relationship of liver xanthine oxidase to quality of, 299.
— effect of varying level and nature of, and level of fat, on development of liver necrosis, 41.
— level and penicillin, effect on growth and intestinal flora of chickens, 175.
— (low-) diets, importance of choline in prevention of nutritional edema in rats fed, 361.
— requirement of rat for maximum growth under conditions of restricted consumption of food, determination of, 133.
Pteroylglutamic acid and leucovorin, comparative biological activity of, 93.
Purified amino acids, growth and metabolism studies with rats fed rations containing, 243.

RALLI, ELAINE P. See Dumm, M. E., 11.
Rat(s), adult, fed diets low in L-, DL- and D-phenylalanine, nitrogen balance of, 437.
— amino acids in production of blood constituents in, 477.
— cotton, occurrence of calcinosis syndrome in. I., 317; II., 327.
— determination of protein requirement of, for maximum growth under conditions of restricted consumption of food, 133.
— effects of a vitamin B_{12} supplement, vitamin B_{12} and streptomycin on metabolism of, 159.
— — — testosterone propionate on niacin metabolism in, 353.
— fed low-protein diets, importance of choline in prevention of nutritional edema in, 361.
— — rations containing purified amino acids, growth and metabolism studies with, 243.
— growing, deposition of fluorine in bones and teeth of, 611.
— immature, fed thyroid, protective effect of dietary fat on, 31.
— infantile, growth-promoting effect of commercial strained meat and fish products investigated with, 119.
— influence of ingested mineral oil upon development of an essential fatty acid deficiency in, 383.
— liver, effects of diet, partial hepatectomy and growth-promoting factors on composition of, 11.
— — fed, administered multiple sublethal doses of x-irradiation, increased survival of, 289.
— tri-o-cresyl phosphate as a vitamin E antagonist for, 583.
— young and adult, with different reserves of vitamin B_{12}, urinary excretion of B_{12} following subcutaneous administration to, 525.
Rations containing purified amino acids, growth and metabolism studies with rats fed, 243.
Reproduction studies with dogs. I. Effect of a corn-soybean meal-alfalfa diet, 621.
Reserves of vitamin B_{12}, urinary excretion of B_{12} following subcutaneous administration to young and adult rats with different, 525.
Riboflavin deficiency in Cebus monkey and its diagnosis, 225.
— diethyl, biological activity of, 539.
— in human serum, 105.
— requirement of baby pig. I. At an environmental temperature of 85°F. and 70% relative humidity, 411.
RICHARDSON, D. See Catron, D. V., 461.
ROACH, GLORIA F. See Westerman, B. D., 147.
ROBBINS, RENEE. See Peterson, D. W., 57.

SALANDER, R. C., AND A. R. PATTON. Free amino acids in chick livers as altered by vitamin B_{12} and fasting, 469.
SANDSTEAD, HAROLD R. See Saunders, J. P., 191.
SARETT, H. P. Effect of oral administration of streptomycin on urinary excretion of B vitamins in man, 275.
SAUNDERS, J. P., H. R. SANDSTEAD, R. E. BUTLER AND O. MICKELSEN. Relation of serum cholinesterase to nutritional status of adolescents, 191.
SCHEID, H. E., M. M. ANDREWS AND B. S. SCHWEIGERT. Comparison of methods for determination of the vitamin B_{12} potency of meats, 601.
SCHWEIGERT, B. S. See Scheid, H. E., 601.
SCOULAR, FLORENCE I. See Harp, M. J., 67.
SEBRELL, W. H., JR., AND E. C. MCDANIEL. Amino acids in the production of blood constituents in rats, 477.
Serum cholinesterase, relation to nutritional status of adolescents, 191.
— human, riboflavin in, 105.
— plasma and white cell-platelet ascorbic acid, daily determination of, in relation to excretion of ascorbic and homogentisic acids by adults maintained on controlled diet, 341.
SHNEOUR, ELIE A. See Peterson, D. W., 57.
SILICIANO, ANN M. See Nasset, E. S., 437.
SIRNY, ROBERT J. See Violante, A. M., 307.
SLINGER, STANLEY J. See Anderson, G. W., 175.
SMITH, A. H. See Lofgreen, G. P., 561.
SNIDER, G. G. See Hoagland, R., 399.
Soybean meal-corn-alfalfa diet, effect of. Reproduction studies with dogs. I., 621.
STARE, FREDRICK J. See Suvarnakich, K., 105.
Sterols (dietary), plasma and liver cholesterol levels, and atherosclerosis, some relationships among, in chicks, 57.
STOKSTAD, EVAN L. R. See Broquist, H. P., 93.
STONE, MIRIAM C. See Westerman, B. D., 147.
STOREVICK, CLARA A. See Davey, B. L., 341.
Strained meat and fish products, commercial, growth-promoting effect of, investigated with infantile rats, 119.
Streptomycin, effect of oral administration on urinary excretion of B vitamins in man, 275.
— vitamin B_{12} and vitamin B_{12} supplement, effects on metabolism of rat, 159.

Subcutaneous administration of vitamin B_{12} to young and adult rats with different reserves, urinary excretion of B_{12} following, 525.
Sublethal doses, multiple, of x-irradiation, increased survival of liver-fed rats administered, 289.
Suckling pigs (see Pigs, suckling).
Survival, increased, of liver-fed rats administered multiple sublethal doses of x-irradiation, 289.
SUVARNAKICH, K., G. V. MANN AND F. J. STARE. Riboflavin in human serum, 105.
SWIFT, C. E. See Hoagland, R., 399.
— RAYMOND W. See Maddy, K. H., 243.
Syndrome, calcinosis, occurrence in cotton rat. I., 317; II., 327.

TEETH and bones of growing rat, deposition of fluorine in, 611.
Temperature, environmental, of 85°F. and 70% relative humidity, riboflavin requirement of baby pig at, 411.
Testosterone propionate, effect on niacin metabolism in rats, 353.
Thiouracil, effect on ascorbic acid requirement of male guinea pig, 487.
Thyroid, protective effect of dietary fat on immature rats fed, 31.
Tract, gastrointestinal (see Gastrointestinal tract).
Tri-o-cresyl phosphate as a vitamin E antagonist for rat and lamb, 583.
Turkey poults, requirement for isoleucine and activities of its isomers for growth of, 631.
Tyrosine content of meat, 307.

UNDERKOFLER, L. A. See Catron, D. V., 461.
Urinary excretion of B vitamins in man, effect of oral administration of streptomycin on, 275.
— — — vitamin B_{12} following subcutaneous administration to young and adult rats with different reserves, 525.

VIOLANTE, A. M., R. J. SIRNY AND C. A. ELVEHJEM. The aspartic acid, glutamic acid, proline and tyrosine content of meat, 307.
Vitamin(s), B, effect of oral administration of streptomycin on urinary excretion in man, 275.
— B_{12} and amino acid imbalances, effect on growth and levels of certain blood constituents in chick, 73.
— — free amino acids in chick livers as altered by, 469.
— — performance of weanling pigs on low levels of, 461.
— — potency of meats, comparison of methods for determination of, 601.
— — requirement of weanling pigs for optimum growth, 461.
— — — — — — — — II. Performance on low levels of vitamin B_{12} and requirement for optimum growth, 461.
— — supplement, vitamin B_{12} and streptomycin, effects on metabolism of rat, 159.
— — urinary excretion folowing subcutaneous administration to young and adult rats with different reserves, 525.
— E antagonist for rat and lamb triocresyl phosphate as a, 583.

WATSON, PATRICIA L. See Mann, G. V., 213; 225.
Weanling pigs (see Pigs, weanling).
Weight (adrenal) and odontoblast height as criteria of ascorbic acid requirement of guinea pig, 503.
WESTERMAN, B. D., G. F. ROACH AND M. C. STONE. Improving the nutritive value of flour. V. The effect of the use of defatted wheat germ, 147.
Wheat germ, defatted, effect of use of. Improving the nutritive value of flour. V., 147.
White cell-platelet, plasma and serum ascorbic acid, daily determination of, in relation to excretion of ascorbic and homogentisic acids by adults maintained on controlled diet, 341.
WIESE, ALVIN C. See Lehrer, W. P., Jr., 203.
WILLIAMS, D. E. See Kratzer, F. H., 631.
— JESSE N., JR. See Bothwell, J. W., 375; Litwack, G., 299.
Women, 6 young, amount of nitrogen retained by, on intake of approximately 70 gm protein a day, 425.
— young college, cobalt metabolism of, on self-selected diets, 67.
WU, MEI-LING. See Davey, B. L., 341.

XANTHINE oxidase (liver), relationship to quality of dietary protein, 299.
X-irradiation, increased survival of liver-fed rats administered multiple sublethal doses of, 289.

YOUNG calf (see Calf, young).
— rats (see Rats, young)
— women (see Women, young).

ZIPKIN, I., AND F. J. MCCLURE. Deposition of fluorine in the bones and teeth of the growing rat, 611.